

Pathological characteristics of uterus in rats with polycystic ovary

Fatemeh Lakzaei¹, Manizheh Karami^{2*}, Maryam Darban Fooladi¹, Mohammadreza Jalali Nadoushan³

1. MSc student, School of Basic Sciences, Shahed University, Tehran, Iran.

2. Associate Professor, School of Basic Sciences, Shahed University, Tehran, Iran.

3. Professor, School of Medicine, Shahed University, Tehran, Iran.

Article info:

Received: 29 April 2013

First Revision: 29 May 2013

Accepted: 20 June 2013

Key Words:

L-Arginine

Polycystic ovary

Pathological evidence

Uterus

A B S T R A C T

Background and Objective: Uterus of rat with polycystic ovary (PCO) may show pathological features. We provided pathological evidence for the rat uterus with NO-induced PCO.

Materials and Methods: Wistar rats (weighing 200-250 g) were kept diestrous to receive L-arginine (50 mg/kg) intraperitoneally (i.p.) for 9 days/once a day. Control group solely received saline (1 ml/kg, 9 days/once per day). At the end of the treatment period, all animals were surgically studied. The rats' ovaries and uteri were examined biometrically and collected in 10% formalin. The pathological data were collectively determined.

Results: The treated ovaries of rats showed polycystic characteristics when compared with the control. The uteri of treated rats also showed pathological changes as compared to those that belonged to the controls.

Conclusion: The pathological aspect of rat uterus may be linked with the cystic characteristic of ovary in PCO model. This study provides pathological evidence for uterus of rat with PCO.

1. Introduction

One of the endocrine disorders of women is known as polycystic ovary (PCO) which is characterized by polycystic feature of ovary along with the hyperandrogenism and ovulatory dysfunction (1,2). Also, one who suffers from PCO may show a higher exhibition of pro-inflammatory agents such as nitric oxide (NO) (3).

We have already shown that the chronic use of L-arginine, a precursor of NO, in rats with diestrous phase may induce the PCO alongside

with lipid metabolism malfunction (3), the characteristics that mark the PCO syndrome (PCOS).

Since, by reviewing of literature, the uteri of those suffering from PCO have not been much studied, we sought to evidence pathologically the PCO model's uterus. In this work, the female Wistar rats were treated chronically with Larginine to provide the PCO model. Also, the experimental animals' uteri were examined to provide valuable data representing the involvement of NO in pathophysiology of the disorder.

***Corresponding Author:**

Manizheh Karami

School of Basic Sciences, Shahed University, Tehran, Iran.

Email: karami@shahed.ac.ir

2. Materials and Methods

2.1. Animals

Wistar rats (body weight 200-250 g) were purchased from Pasteur Institute of Iran and were retained under standard conditions ($21 \pm 3^\circ\text{C}$ and 12-h light/dark cycle) with food and water ad libitum. All experiments were approved by local Ethical committee.

2.2. Drugs

L-arginine (Merck Co., Germany) was injected i.p. for a 9-day period/ once per day. The vehicle (saline at 1 ml/kg, i.p.) was used in control group.

2.3. Female Cycle Test

Since female rats with 4-5 day sexual cycle are always in diestrus (4,5) unless in case of having mating with the male rat, so, the rats were kept virgin in the present study to avoid the change in female sexual cycle.

2.4. Drug administrations

Animals were randomly divided into the L-arginine (50 mg/kg), and saline control (1 ml/kg) groups (n=6). They were injected the agent or saline intraperitoneally (i.p.) once a day during the 9 days period.

2.5. Surgery procedure

The treatment groups were anesthetized by an overdose of diethyl ether. Then, a midline incision in the lower abdomen area was performed. The ovaries and uteri were biometrically examined and dissected out. They were collected in 10% formalin for histological examination.

2.6. Histological investigation

The collected tissues were processed and sectioned at a thickness of 3-4 μm . They were stained by Hematoxylin and Eosin (H&E) method (6). The thin sections were then dehydrated, cleared, and eventually mounted with entellane (Merck Co., Germany) and coverslipped. The prepared slides were evaluated with light microscope (Olympus, Japan) at 4-40X.

2.7. Image analysis

The photomicrographs were assessed in areas of $100\text{-}\mu\text{m}^2$ with an aid of Image Tool program (UTHSCSA, version 2.03), the free image processing and analysis program for Microsoft Windows.

2.8. Statistical analysis

All data were first assessed by Kolmogorov-Smirnov (K-S) to show the equality to analysis by variance (ANOVA). The ANOVA was then performed using SPSS software (version 13.0; SPSS, Inc., Chicago, IL), followed by post-hoc test. Statistical significance was considered at $p < 0.05$. All data are expressed as Means \pm SEM. The photos were examined in an area of $100\text{-}\mu\text{m}^2$ using the Image Tool program.

3. Results

3.1. Histology

The ovaries from the L-arginine-treated group (50 mg/kg, chronically) showed cystic formations (Fig. 1B) as compared to control samples (Fig. 1A), the aspects certifying the polycystic ovary (PCO) example.

The uteri samples of those received chronically L-arginine (50 mg/kg) (Fig. 1C) presented pathological evidence to those belonged to the controls (Fig. 1D). Due to activation of the L-arginine-related metabolic pathway, the uterus wall revealed the aspects of swelling, the proliferation and angiogenesis, suggesting an inflammatory process involvement.

3.2. Biometrical value (diameters of uteri)

The uteri diameters were calculated in all groups. They showed changes in L-arginine-treated group as compared to those obtained from saline group. The uteri of rats treated with L-arginine showed a significant increase when compared with the control group ($p < 0.05$) (Fig.2).

4. Discussion

This research study showed that chronic treatment of rats with a nitric oxide (NO) agent, L-arginine, induces the polycystic ovary (PCO) formation as well as the uteri inflammation.




Figure 1. Photomicrographs of ovaries and uteri from control (A, C), and L-arginine-treated (B, D) rats.




Figure 2. The diameter of uteri in rats. X axis denotes the control and experimental groups (n=6). Control was injected saline (1 ml/kg, i.p., 9 days/once per day). The experimental rats received L-arginine (50 mg/kg, i.p. for 9 days). Values are means \pm S.E.M.* $p < 0.05$ vs. control (based on *Post-hoc* test).

The pro-inflammatory NO participates in endocrine physiological and pathophysiological events (7). The enzyme nitric oxide synthase (NOS) produces the NO by the oxidation of terminal guanidino nitrogen of arginine (7). We have already demonstrated the NOS activation with NADPH-diaphorase (8). The molecule NO is well introduced as a local inflammatory generator (9). The present work also provides support for a functional role of the NO in the ovarian and uterine events. This plan further supports our previous study (8), denoting that hyperactivity of enzyme NOS due to chronic usage of L-arginine induces polycystic formation in treated rats' ovaries. In agreement with this idea, it has been indicated that the presence of large cysts due to treatment by NO producer, L-arginine, accords with common characteristics of PCOS (10).

In addition of significant changes in feature of ovary, the uterus of the L-arginine-treated rats also showed differences as compared to the saline control group. These finding may suggest that the NO as a pro-inflammatory element may induce the inflammatory changes in uteri as well as ovaries. Although the exact effect of the L-arginine in this study remained elusive, the metabolic pathway may involve the NO which is known as a short-lived cytotoxic mediator (11). By viewing of the uterus diameter that increased in L-arginine-treated rats, it appears that inflammatory processes play crucial role in reproduction at all levels from the follicles and ovarian function to the accessory sex organs (i.e. uterus). We aimed to involve the NO by chronic use of L-arginine in PCO and pathology of rats' uteri. We have now evidenced that production of pro-inflammatory NO may induce significant change in the ovary and uterus parameters. It should be notified that role of NO in activation of NOergic neurons of the pelvic plexus has been previously shown (12). The NO has also been involved in the control of uterine smooth muscle via NOergic terminals (13). Based on our results, however, the exact mechanisms to involve the inflammatory processes in ovary and uterus events rest elusive.

In conclusion, this study indicates the pathological evidence in reproductive system of Wistar rats with PCO. Because of activation of the L-arginine-related metabolic pathway, the uterus wall illustrated the aspects of swelling, the

proliferation and angiogenesis, suggesting an inflammatory process involvement.

Acknowledgments

The authors are thankful to Mr Vahid Yeghaneh Kaffash for expert assistance in histology procedure. We also wish to thank Research Deputy at Shahed University and Neurophysiology Research Center of Shahed University.

Conflict of interest

The authors state that they do not have any conflict of interest.

References

- 1-Amowitz LL, Sobel BE. Cardiovascular consequences of polycystic ovary syndrome. *Endocrinology and Metabolism Clinics of North America* 1999; 28: 439-458.
- 2-Dokras A. Cardiovascular disease risk factors in polycystic ovary syndrome. *Seminars in Reproductive Medicine* 2008; 26: 39-44.
3. Apridonidze T, Essah PA, Iuorno MJ, Nestler JE. Prevalence and characteristics of the metabolic syndrome in women with polycystic ovary syndrome. *The Journal of Clinical Endocrinology and Metabolism* 2005; 90: 1929-35.
4. Maeda KI, Kura SO, Tsukamura H. Physiology of reproduction, in the laboratory Rat: the handbook of experimental animal. Hrinke CJ, Editor. London: Academic press 2000; 145-176.
5. Mohammadzadeh A, Heidari M, Soltan Ghoraii H, Zarnani AH. Induction of endometriosis by implantation of endometrial fragments in female rats. *Iranian Journal of Reproductive Medicine* 2006; 4: 63-67.
6. Manneras L, Cajander S, Holmang A, Seleskovic Z, Lystig T, Lonn M, Stener-Victorin E. A new rat model exhibiting both ovarian and metabolic characteristics of polycystic ovary syndrome. *Endocrinology* 2007; 148: 3781-3791.
7. Zackrisson U, Mikuni M, Wallin A, Delbro D, Hedin L, Brännström M. Cell-specific localization of nitric oxide synthases (NOS) in the rat ovary during follicular development, ovulation and luteal formation. *Human Reproduction* 1996; 11: 2667-2673.

8. Hassani F, Karami M, Jalali Nadoushan MR, Eftekhari Yazdi P. Nitric oxide-induced polycystic ovaries in the Wistar rat. *International Journal of Fertility and Sterility* 2012; 6: 111-116.
9. Nakamura Y, Kashida S, Nakata M, Takiguchi S, Yamagata Y, Takayama H, and et al. Changes in nitric oxide synthase activity in the ovary of gonadotropin treated rats: The role of nitric oxide during ovulation. *Endocrine Journal* 1999; 46: 529-538.
10. Legro RS. Polycystic ovary syndrome: the new millenium. *Molecular and Cellular Endocrinology* 2001; 184: 87-93.
11. Drapier JC, Wietzerbin J, Hibbs JB Jr. Interferon- γ and tumor necrosis factor induce the l-arginine-dependent cytotoxic effector mechanism in murine macrophages. *European Journal of Immunology* 1988; 18: 1587-1592.
12. Burnett AL, Lowenstein CJ, Brecht DS, Chang TSK, Snyder SH. NO: A physiologic mediator of penile erection. *Science* 1992; 257: 401-403.
13. Brännström M, Janson PO. The role of leukocytes and cytokines as paracrine regulators in the mechanisms of ovulation. In: Fujimoto S, Hsueh AJW, Strauss IH, editors. *Frontiers in Endocrinol.* Vol 13. New York: Ares-Serono Symposium Pub; 1995; 225-233.