



Antifungal effect of *Heracleum persicum* on standard *Candida dublinensis* strain: An in Vitro study

Fatemeh Azizi¹, Sayed Shojaedin Shayegh², Mohammad Amin Bafandeh², Alireza Naeini^{3*}

1. Faculty of Dentistry, Shahed University, Tehran, Iran

2. Department of Prosthodontics, Faculty of Dentistry, Shahed University, Tehran, Iran

3. Department of Parasitology and Mycology, Faculty of Medicine, Shahed University, Tehran, Iran

Abstract

Background and Objective: The aim of this study was to investigate the antifungal effects of aqueous, alcoholic, acetone extracts, and essential oil of *Heracleum persicum* seeds (*Golpar*) against standard *Candida dublinensis* (*C.d*) strain.

Materials and Methods: The antifungal effects were evaluated using the agar well diffusion method. Additionally, the *Minimum Inhibitory Concentration* (MIC) and *Minimum Fungicidal Concentration* (MFC) were determined using the serial dilution method. Standard strain *C. d* was investigated, and Nystatin was used as the positive control.

Results: The mean diameter of growth inhibition zones for the alcoholic and acetone extracts and essential oil of *Golpar* seeds were reported in millimeters as follows: 13 mm and 10 mm and 15 mm for *C.d*, respectively. Based on the results, the MIC and MFC of the alcoholic extract of *Golpar* seeds against *C.d* were reported as 40 mg/ml and 40 mg/ml. Additionally, the MIC and MFC of the acetone extract of *Golpar* seeds against *C. d* were reported as 14 mg/ml and 28 mg/ml, respectively. Moreover, MIC of the essential oil of *Golpar* seeds against *C.d* was 0.5 mg/ml and the MFC was 1 mg/ml.

Conclusion: According to the results, the essential oil and the alcoholic and acetone extracts of *Golpar* seeds demonstrated promising antifungal and anti- *Candida* activity against *C.d*. Therefore, based on the findings of this study and further research on the extracts and essential oil of *Golpar*, their antifungal and anti-*Candida* effects could be utilized in the formulation of mouthwashes for patients with removable dentures suffering from Oral Candidiasis.

Keywords: *Heracleum persicum*, *Golpar*, *C. dublinensis*, Mouthwash, Removable Dentures

1. Introduction

The use of complete removable dentures poses challenges for both patients and clinicians. Issues arising from the use of complete dentures for patients and clinicians, particularly in relation to oral hygiene neglect leading to microbial proliferation in the denture, are highlighted as the main cause of denture stomatitis (1, 2). This introduction also addresses the prevalence of denture stomatitis in Iran and the contributing factors to this condition (3, 4). Various treatments, including antifungal therapies and denture replacement or adjustment, are examined for

controlling the symptoms of denture stomatitis (5, 6, 7).

This section of the article explores the main problem associated with denture stomatitis and its causative factors (3, 8). Among the causative factors of denture stomatitis are candidal infections, particularly with *Candida* species. Various factors such as poor denture hygiene, improper use of antifungal medications, and compromised immune system can lead to the development of denture stomatitis (9, 10). It has been investigated that various species of *Candida*, especially *Candida albicans*, are the primary cause of denture stomatitis (3, 11, 12, 13). Additionally, other

species of *Candida* such as *Candida glabrata* and *Candida dubliniensis* are also discussed (14). These species may exhibit greater resistance to antifungal treatments (13, 15).

Furthermore, this research examines the antimicrobial and antifungal effects of the extract and essential oil of *Heracleum persicum* (commonly known as Golpar). This plant contains compounds that can effectively control the growth of fungal species, especially *Candida*. Additionally, this plant possesses antioxidant and immunomodulatory properties that could be effective in inhibiting and preventing the occurrence of denture stomatitis (16, 17, 18).

Given the limited information available regarding the efficacy of Golpar extracts and essential oil in inhibiting and controlling the growth of fungal species, further laboratory studies are proposed to evaluate the antifungal properties of this plant and its use as a therapeutic option for denture stomatitis.

In summary, this study focuses on the antifungal effects of Golpar seed extracts and essential oil on different fungal species and emphasizes their importance in controlling denture stomatitis and reducing resistance to chemical antifungal agents. This research introduces innovation in examining the different effects of herbal products on fungi and the impact of processing variations on their properties.

This disease poses many challenges for patients and clinicians using complete removable dentures (3). Many symptoms and factors, including poor oral hygiene and continuous denture use, are associated with denture stomatitis (19). Treating this disease requires a comprehensive treatment plan and identifying all predisposing factors (20).

Candida dubliniensis:

C. dubliniensis is an opportunistic pathogen first identified in AIDS patients (21). This fungus is commonly found alongside *C. albicans* in immunocompromised patients but differs morphologically (22). Cases of *C. dubliniensis* identification in oral samples of diabetic patients or those using insulin have been reported (23). *C. dubliniensis* is sensitive to fluconazole but resistance to this drug has also been reported (24).

Heracleum persicum (Golpar):

Heracleum persicum, commonly known as Golpar, is a member of the Apiaceae family and is native to Iran. Its seeds have long been used as a spice, and its young blossoms in pickles and traditional medicine. Important furanocoumarins extracted from the roots of this plant include Pimpinellin, Isopimpinellin, bergapten, isobergapten, and sphonodin. The main components of Golpar seed extracts include hexyl butyrate, octyl acetate, hexyl 2-methyl butanoate, and hexyl isobutyrate (25, 26). This plant has various

properties in the field of medicine and traditional medicine, including inhibiting oxidative stress processes and possessing antioxidant properties (27, 28, 29, 30). Additionally, its anti-inflammatory and antimicrobial properties are noteworthy (31, 32). Some terpene compounds and alkaloids present in Golpar exhibit anticonvulsant and analgesic properties (33, 34). The use of Golpar seeds can help stimulate the human humoral and cellular immune systems, but attention should be paid to the amount of consumption and its limitations (35).

2. Materials and Methods

2.1. Research method

This study was conducted as a laboratory experiment using the well diffusion and macrobroth dilution methods (serial dilution method) to investigate the anti-candidal effect of Golpar seed alcohol and acetone extracts (36).

2.2. Preparation of essential oil

The essential oil was prepared using the water distillation method. To prepare the oil, 100 grams of dried Golpar seed petals were ground using an electric grinder and placed in the balloon of the distillation device. Then, 700 milliliters of distilled water was added to the balloon, and the distillation process continued for two hours. Finally, the collected oil volume was determined and stored in the refrigerator for preservation (37).

2.3. Preparation of extract

Extraction was performed using the immersion method (maceration) with ethanol and acetone solvents. A quantity of 100 grams of the plant material, after grinding, was placed in extraction vessels, and 100% acetone and 80% ethanol solvents were added. The solutions were then stored at laboratory temperature for 3 to 4 days (37).

2.4. Preparation of aqueous extract

The Seed of plant (100 g) was cleaned and powdered using a mechanical grinder. The powder (100 g) was added to 400 mL hot water, boiled for 15 minutes and filtered through a Whatman paper (No. 42). The filtrate was evaporated to dryness under reduced pressure to obtain a viscous residue. The residue was suspended in normal saline (37). It is necessary to mention that the plant extract solutions were sterile.

2.5. Well diffusion assay

In this assay, wells with a diameter of approximately 6 millimeters were created on Sabouraud dextrose agar plates, and the experimental solutions of Golpar seed extracts were poured into the wells. After 24 to 48 hours, areas of fungal growth inhibition were

examined. Finally, the macrobroth dilution assay steps were completed (37).

2.6. Study population

Standard strain *C. dubliniensis* ATCC = CD60 was selected for the study.

2.5. Statistical analysis

Using SPSS software and One-Way ANOVA test followed by Tukey's post-hoc test, the results were analyzed. The tests indicated that the results of the experiments on Golpar seed water extract were ineffective and were not included in the statistical analysis.

3. Results

3.1. Results on *C. dubliniensis* strain

Investigation showed that the diameter of the growth inhibition zone of Golpar seed essential oil was significantly higher compared to its acetone extract. However, no significant difference was observed compared to its ethanol extract. Furthermore, the diameter of the growth inhibition zone for both extract and essential oil was significantly lower than nystatin. This was also true for MIC and MFC.

These results demonstrate that Golpar seed extracts and essential oil can be considered as effective options in controlling candidal pathogens.

4. Discussion

Candida species are part of the normal flora of the oral cavity, but under conditions such as immunodeficiency and malnutrition, they can cause opportunistic superficial skin and mucosal infections. *C. albicans* is the main causative agent of oral candidiasis, but nowadays, other Candida species are also observed as causative agents. In this study, we investigated the antifungal effects of aqueous, ethanol, and acetone extracts, as well as the essential oil of Golpar seeds against *C. dubliniensis*.

Golpar seed has been traditionally used as an anti-inflammatory and analgesic drug in traditional medicine and its antioxidant and antimicrobial effects have been recognized in laboratory settings. Studies have shown that Golpar seed essential oil has moderate antifungal effects (38). Additionally, in other studies, the inhibitory effect of Golpar seed essential oil on the growth of *Candida Zeylonoides* has been reported (39).

In the present study, the antifungal effect of Golpar seed essential oil in inhibiting the growth of *C. dubliniensis* fungi was greater than its extract form. The stronger activity of essential oils in inhibiting fungal growth may be due to the synergistic effects of active components of the essential oil compared to the individual compounds that make up the extract (40, 41).

The findings of these investigations have shown that

the aqueous Golpar extract due to its abundant polar groups ionizes the surrounding environment of the fungus, minimizing the interference of fungal compounds with the surrounding environment, which does not have a measurable effect on the growth inhibition of *C. dubliniensis* (42). According to studies, *C. dubliniensis* have shown the highest resistance to Golpar seed extracts, which is consistent with our findings. Additionally, the fungal species *C. dubliniensis* quickly develops resistance to antifungal drugs and compounds, which may be due to its ability to survive in harsh environmental conditions (43).

In other research, it has been shown that the antioxidant compounds present in Golpar plant stimulate the proliferation of B lymphocytes and macrophages, playing an important role in strengthening the humoral and cellular immune system (44). Golpar seeds contain a variety of compounds including alkaloids, triterpenes, and terpenoids responsible for the plant's anticonvulsant effects (45, 46).

Conclusion

Based on the findings of the present study, it can be concluded that the seed extract of the Golpar plant can be used effectively in combating candida species in denture stomatitis infections, given its efficacy comparable to that of nystatin. Moreover, the efficacy of both alcoholic and acetone forms of Golpar seed extract against *C. dubliniensis* is similar, and the choice between these two can be made based on other parameters such as side effects and cost-effectiveness.

Limitations and recommendations

One of the limitations of this study is its laboratory-based nature. The growth and colonization conditions of microbial species in the oral cavity and under dentures may differ from laboratory conditions. Although denture stomatitis-causing species are highly diverse, only four species were investigated in this study. Future studies are recommended to explore more parameters in this regard.

Since most studies have focused on *C. albicans* strains and the mechanism of action of Galper extract has been examined, it is suggested to investigate the effects of these plant derivatives on non-albicans strains in further studies. Typically, the efficacy of extracts and essences in converting these derivatives into mouthwashes and their interaction with other auxiliary substances or existing oxidants in mouthwashes may vary. Clinical trials in the future are recommended to examine the antifungal effects of the mentioned compounds in the form of mouthwashes under controlled conditions.

References

- White DA, Tsakos G, Pitts NB, Fuller E, Douglas GV, Murray JJ, et al. Adult Dental Health Survey 2009: common oral health conditions and their impact on the population. *British Dental Journal* 2012; 213(11):567-72.
- Coulthwaite L, Verran J. Potential pathogenic aspects of denture plaque. *British Journal of Biomedical Science*. 2007; 64(4):180-9.
- Moosazadeh M, Akbari M, Tabrizi R, Ghorbani A, Golkari A, Banakar M, et al . Denture Stomatitis and *Candida albicans* in Iranian Population: A Systematic Review and MetaAnalysis. *Journal Dental (Shiraz)*. 2016; 17(3 supply):283-292.
- Scully C, Porter S. ABC of oral health. Swellings and red, white, and pigmented lesions. *British Medical Journal* 2000; 321: 225–228.
- Akpan A, Morgan R. Oral candidiasis. *Postgraduate Medicine Journal*. 2002; 78(922):455-9.
- Martorano-Fernandes L, Dornelas-Figueira LM, Marcello-Machado RM, Silva RB, Magno MB, Maia LC et al . Oral candidiasis and denture stomatitis in diabetic patients: Systematic review and meta-analysis. *Brazilian oral research journal*. 2020, 21; 34:e113.
- Salerno C, Pascale M, Contaldo M, Esposito V, Busciolano M, Milillo L, et al. *Candida*-associated denture stomatitis. *Medicina Oral Patologia Cirurgia Buca*. 2011, 1; 16(2):e139-43.
- Galvan R, McBride M, Koriath TV, Garcia-Godoy F. Denture Hygiene as It Relates to Denture Stomatitis: A Review. *Compendium of Continuing Education in Dentistry* . 2021; 42(4):e1-e4.
- Contaldo M, Romano A, Mascitti M, Fiori F, Della Vella F, Serpico R, et al. Association between denture stomatitis, *Candida* species and diabetic status. *Journal of Biological Regulators and Homeostatic Agents*. 2019; 33(3 Suppl. 1):35-41.
- Yano J, Yu A, Fidel PL Jr, Noverr MC. *Candida glabrata* Has No Enhancing Role in the Pathogenesis of *Candida*-Associated Denture Stomatitis in a Rat Model. *mSphere*. 2019 3; 4(2):e00191-19.
- Silva S., Henriques M., Hayes A., Oliveira R., Azeredo J., Williams D.W. *Candida glabrata* and *Candida albicans* co-infection of an in vitro oral epithelium. *Journal of Oral Pathology Medicine*. 2011; 40:421–427.
- Manikandan S, Vinesh E, Selvi DT, Kannan RK, Jayakumar A, Dinakaran J. Prevalence of *Candida* among Denture Wearers and No denture Wearers. *Journal of Pharmacy and Bio allied Sciences*. 2022; 14(Supply 1):S702-S705.
- Ferreira R, Limeta A, Nielsen J. Tackling Cancer with Yeast-Based Technologies. *Trends Biotechnol*. 2019; 37(6):592-603.
- Maluche M.E., Santos J.I. *Candida* spp e infecções hospitalares: aspectos epidemiológicos e laboratoriais. *RBAC*. 2008; 40:65–67.
- Coco, B.J., Bagg, J., Cross, L.J., Jose, A., Cross, J. and Ramage, G., Mixed *Candida albicans* and *Candida glabrata* populations associated with the pathogenesis of denture stomatitis. *Oral Microbiology and Immunology*, 2008 23: 377-383.
- Johnson, John A., Duncan Webster, and Christopher A. Gray. "The Canadian medicinal plant *Heracleum maximum* contains antimycobacterial diynes and furanocoumarins." *Journal of ethnopharmacology* 2013: 232-237.
- Nejad, Batool S., Rajabi, Mahsa, Mamoudabadi, Ali Z., and Majid Zarrin. "In Vitro Anti-*Candida* Activity of the Hydroalcoholic Extracts of *Heracleum persicum* Fruit against Pathogenic *Candida* Species." *Jundishapur Journal of Microbiology*. 2014 January; 7(1): e8703.
- Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nature Protocols*. 2008; 3(2):163-75.
- Gual-Vaqués P., Jané-Salas E., Egido-Moreno S., Ayuso-Montero R., MariRoig A., López-López J. Inflammatory papillary hyperplasia: a systematic review. *Medicina Oral, Patologia Oral Y Cirugia Bucal* . 2017; 22(1):e36–e42.
- Sampaio-Maia B., Figueiral M. H., Sousa-Rodrigues P., Fernandes M. H., Scully C. The effect of denture adhesives on *Candida albicans* growth in vitro. *Gerodontology*. 2012; 29(2):e348–e356.
- Pristov KE, Ghannoum MA. Resistance of *Candida* to azoles and echinocandins worldwide. *Clinical Microbiology Infection*. 2019; 25(7):792-798.
- Polke M, Hube B, Jacobsen ID. *Candida* survival strategies. *Advance Apply Microbiology*. 2015; 91:139-235.
- Jabri B, Iken M, Ait-Ou-Amar S, Rida S, Bouziane A, Ennibi OK. *Candida albicans* and *Candida dubliniensis* in Periodontitis in Adolescents and Young Adults. *International Journal Microbiology*. 2022 11; 2022:4625368.
- Petty LA, Gallan AJ, Detrick JA, Ridgway JP, Mueller J, Pisano J. *Candida dubliniensis* Pneumonia: A Case Report and Review of Literature. *Mycopathologia*. 2016; 181(9-10):765-8.
- Majidi Z, Lamardi SS. Phytochemistry and biological activities of *Heracleum persicum*: A review. *Journal of integrative medicine*. 2018 1; 16(4): 223- 35. C

26. Sefidkon F, Dabiri M, Mohammad N. Analysis of the oil of *Heracleum persicum* L.(leaves and flowers). *Journal of Essential Oil Research*. 2002 1; 14(4):295-7. E
27. Changxin g L, Dongfang D, Lixue Z, Saeed M, Alagawany M, Farag MR, Chenling M, Jianhua L. *Heracleum persicum*: Chemical composition, biological activities and potential uses in poultry nutrition. *World's Poultry Science Journal*. 2019 1; 75(2):207-18. I
28. Asgarpan ah J, Dadashzadeh Mehrabani G, Ahmadi M, Ranjbar R, Safi-Aldin Ardebily M. Chemistry, pharmacology and medicinal properties of *Heracleum persicum* Desf. Ex Fischer: A review. *Journal of Medicine Plants Research*. 2012 16; 6(10):1813-20. L
29. Sharififar F, Pournourmohammadi SH, Arabnezhad M, Rastegarianzadeh R, Ranjbaran O, Pourhemati A. Immunomodulatory activity of aqueous extract of *Heracleum persicum* Desf. in mice.
30. Mojab F, Nikavar B. Composition of the Essential Oil of the Root of *Heracleum persicum* from Iran. 2011
31. Sayyah M, Moaied S, Kamalinejad M. Anticonvulsant activity of *Heracleum persicum* seed. *Journal of ethnopharmacology*. 2005 8;98(1-2):209-11.
32. Hoseinifara SH, Zoheiri F, Lazado CC. Dietary phytoimmunostimulant Persian hogweed (*Heracleum persicum*) has more remarkable impacts on skin mucus than on serum in common carp (*Cyprinus carpio*). *Fish & shellfish immunology*. 2016 1; 59:77-82.
33. Coruh NU, Celep AS, Özgökçe F. Antioxidant properties of *Prangos ferulacea* (L.) Lindl., *Chaerophyllum macropodium* Boiss. and *Heracleum persicum* Desf. from Apiaceae family used as food in Eastern Anatolia and their inhibitory effects on glutathione-S-transferase. *Food chemistry*. 2007 1; 100(3):1237-42.
34. Davari M, Ezazi R. Chemical composition and antifungal activity of the essential oil of *Zhumeria majdae*, *Heracleum persicum* and *Eucalyptus* sp. against some important phytopathogenic fungi. *Journal de mycologie medicale*. 2017 1; 27(4):463-8.
35. Fazeli M, Alizadeh M, Pirsas S. Nanocomposite film based on gluten modified with *Heracleum persicum* Essence/MgO/Polypyrrole: investigation of physicochemical and electrical properties. *Journal of Polymers and the Environment*. 2022; 30(3):954-70.
36. Naeini, A., Yaraee, R., Shokri, H. Antifungal and immunomodulatory activity of *Allium jesdianum* Boiss extracts. *Journal of Herbal Medicine pharmacology*. 2020;9(1):75-80
37. Naeini A, Naseri M, Kamalinejad M, Khoshzaban F, Rajabian T, Nami H, et al . Study on Anti_candida Effects of Essential Oil and Extracts of Iranian Medicinal Plants, In vitro. *Journal of Medicine Plants* 2011; 10 (38):163-172.
38. MimicaDukic N, Bozin B, Sokovic M, Mihajlovic B, Matavulj M. Antimicrobial and antioxidant activities of three *Mentha* species essential oils. *Planta Medicine*. 2003;69(5):413–9.
39. Shokri A, Sharifzadeh A, Ashrafi L. Anti-Candida zeylanoides activity of some Iranian plants used in traditional medicine. *Journal of Medical Mycology*. 2012; 22(2):211- 216.
40. harifzade h A, Shokri H, Katirae F. Anti-adherence and Anti-fungal Abilities of Thymol and Carvacrol Against Candida Species Isolated From Patients with Oral Candidiasis in Comparison with Fluconazole and Voriconazole. *Jundishapur Journal of Natural Pharmaceutical Products*. 2021; 16(1):e65005.
41. Mostafa AA, Yasin MT, Al-Askar AA, Al-Otibi F. Phytochemical analysis, antiproliferative and antifungal activities of different *Syzygium aromaticum* solvent extracts. *Journal of King Saud University - Science*. 2023;35(1):102362
42. Yarborough A, Cooper L, Duqum I, Mendonça G, McGraw K, Stoner L. Evidence Regarding the Treatment of Denture Stomatitis. *Journal of Prosthodontology*. 2016; 25(4):288-301.
43. Hemati A, Azarnia M, Angaji SA. Medicinal effects of *Heracleum persicum* (Golpar). *Inflammation*. 2010 1; 9:10.
44. Mustafavi SH, Abbasi A, Morshedloo MR, Pateiro M, Lorenzo JM. Essential Oil Variability in Iranian Populations of *Heracleum persicum* Desf. ex Fischer: A Rich Source of Hexyl Butyrate and Octyl Acetate. *Molecules*. 2022 23;27(19):6296.
45. Sudbery P, Gow N, Berman J. The distinct morphogenic states of *Candida albicans*. *Trends Microbiology*. 2004; 12:317–24.
46. Staib P, Morschhäuser J. Chlamydospore formation in *Candida albicans* and *Candida dubliniensis*--an enigmatic developmental programme. *Mycoses*. 2007; 50:1–12.